

Sub
B
AI 2. (Amended) The method according to claim 11, wherein the intermittent administration of Org 33245 takes place as an addition to progestagen-only therapy.

Sub
B
AI 4. (Amended) The method of claim 11, wherein Org 33245 is administered for 1-7 days during a cycle of 28-32 days, wherein one dosage marks the end of a cycle and the optional other dosages are administered regularly divided over the remaining days of the cycle.

AI 5. (Amended) A contraceptive kit for the daily administration of a progestagen and for the intermittent administration of an anti-progestagen, comprising a progestagen dosage unit and an anti-progestagen dosage unit, wherein the anti-progestagen dosage unit comprises Org 33245.


6. (Amended) A combined dosage unit comprising a progestagen and an anti-progestagen, wherein the anti-progestagen is Org 33245.

7. (Amended) A method of contraception comprising daily administering to a female of childbearing age a contraceptively effective amount of a progestagen and intermittently administering an anti-progestagen, wherein the anti-progestagen is Org 33245.

8. (Amended) A method of treatment of irregular or breakthrough uterine bleeding in a female using a progestagen-only preparation, comprising intermittently administering an anti-progestagen, wherein the anti-progestagen is Org 33245.

9. (Amended) The method according to claim 7, wherein the anti-progestagen is administered on 1-4 days in a cycle of 28-32 days.


-- 10. A method of anti-progestagen therapy, comprising administering an effective amount of the anti-progestagen of Formula II



Formula II

wherein X is selected from the group consisting of (H,H), (O) and (N-OH), or a salt thereof, said anti-progestagen being administered intermittently, whereby the time between each pair of sequentially administered doses of anti-progestagen is greater than one day. --

-- 11. A method of anti-progestagen therapy, comprising administering an effective amount of the anti-progestagen Org 33245



Org 33245

or a salt thereof, said anti-progestagen being administered intermittently, whereby the time between each pair of sequentially administered doses of anti-progestagen is greater than one day. --

-- 12. The method of claim 11, wherein the anti-progestagen therapy is hormone replacement therapy. --

Sub
B1
-- 13. The method claim 11, wherein the anti-progestagen therapy is contraception. --

-- 14. The method of claim 11, wherein the anti-progestagen therapy is for minimizing uterine bleeding. --

173
-- 15. The method according to claim 8, wherein the anti-progestagen is administered on 1-4 days in a cycle of 28-32 days. --

-- 16. The method of claim 10, wherein the anti-progestagen therapy is hormone replacement therapy. --

Sub
B1
-- 17. The method of claim 10, wherein the anti-progestagen therapy is contraception. --

-- 18. The method of claim 10, wherein the anti-progestagen therapy is for minimizing uterine bleeding. --